CONTENTS

VOLUME 500

Volume 500
by M. Atkins 1

The *Journal of Chromatography* family album
by M. Lederer 3

Hypercoulometry of the second kind in electron-capture detection
by W. A. Aue (Halifax, Canada) and K. W. M. Siu, D. Beauchemin and S. S. Berman (Ottawa, Canada) (Received September 20th, 1989) 95

Organic solutes in paraffin solvents. Influence of the size of the solvent molecule on solution data
by G. Défayes, D. F. Fritz, T. Görner, G. Huber, C. de Reyff and E. sz. Kováts (Lausanne, Switzerland) (Received September 4th, 1989) 139

Theoretical analysis of non-linear preparative liquid chromatography
by B. Lin, Z. Ma, S. Golshan-Shirazi and G. Guiochon (Oak Ridge, TN, U.S.A.) (Received June 19th, 1989) 185

Limiting activity coefficients and gas–liquid partition coefficients of alkylbenzenes in hydro–organic solvents
by W. J. Cheong and P. W. Carr (Minneapolis, MN, U.S.A.) (Received August 18th, 1989) 215

Diffusion coefficients in water–acetonitrile mixtures
by V. Huss, J. L. Chevalier and A. M. Siouffi (Marseille, France) (Received July 20th, 1989) 241

Selection of the solid support — a crucial step for the realization of solvent generated liquid–liquid chromatography
by J. F. K. Huber (Vienna, Austria), M. Pawlowska (Warsaw, Poland) and P. Markl (Vienna, Austria) (Received October 20th, 1989) 257

Possibilities of determination and prediction of solute capacity factors in reversed-phase systems with pure water as the mobile phase
by P. Jandera and J. Kubát (Pardubice, Czechoslovakia) (Received October 16th, 1989) . 281

Modelling of cation retention in ion chromatography using fixed-site and dynamically coated ion-exchange columns
by P. R. Haddad and R. C. Foley (Kensington, Australia) (Received September 13th, 1989) 301

Further studies of diethyldithiocarbamate complexes by high-performance liquid chromatography
by S. Dilli, P. R. Haddad and A. K. Htoon (Kensington, Australia) (Received August 9th, 1989) 313

Experimental protocol for the assessment of solvent strength and selectivity of liquid phases used in gas chromatography
by S. K. Poole and C. F. Poole (Detroit, MI, U.S.A.) (Received October 4th, 1989) . . . 329

Method for radio-capillary gas chromatography employing a modified oxidation–reduction train and flow-through detector
by R. B. Croteau and D. M. Satterwhite (Pullman, WA, U.S.A.) (Received September 7th, 1989) 349

Gas–liquid chromatography in qualitative analysis. XVIII. Study of the dependence of the retention behaviour of polyoxyethylene glycol columns on the nature of the support and phase loading
by J. J. Dowling and M. B. Evans (Hatfield, U.K.) and J. K. Haken (Kensington, Australia) (Received August 30th, 1989) 355

Analysis of polycarbonate–polydimethylsiloxane block copolymers
by J. K. Haken, N. Harahap and R. P. Burford (Kensington, Australia) (Received July 20th, 1989) 367

Selection of electrophoric derivatives of 1-aminopyrene and 2-aminofluorene for determination by gas chromatography with electron-capture negative-ion mass spectrometry
by J. Bakthavachalam and R. S. Annan (Boston, MA, U.S.A.), F. A. Beland (Jefferson, AR, U.S.A.) and P. Vouros and R. W. Giese (Boston, MA, U.S.A.) (Received September 25th, 1989) 373

Use of stable isotopes and gas chromatography–mass spectrometry in the study of different pools of neurotransmitter amino acids in brain slices
by I. M. Kapetanovic, W. D. Yonekawa and H. J. Kupferberg (Bethesda, MD, U.S.A.) (Received August 1st, 1989) 387

Opioid and tachykinin peptides, and their precursors and precursor-processing enzymes, in human cerebrospinal fluid
by D. Liu, C. Dass, G. Wood and D. M. Desiderio (Memphis, TN, U.S.A.) (Received October 11th, 1989) 395

Separation and quantification by gas chromatography–mass spectrometry of arabinitol enantiomers to aid the differential diagnosis of disseminated candidiasis
by J. Roboz, E. Nieves and J. F. Holland (New York, NY, U.S.A.) (Received August 14th, 1989) 413

Gas–liquid chromatographic profiling of plasma lipids using high-temperature-polarizable capillary columns
by A. Kuksis and J. J. Myher (Toronto, Canada) and P. Sandra (Ghent, Belgium) (Received October 16th, 1989) 427

Automated high-performance liquid chromatographic determination of plasma free fatty acids using on-line derivatization with 9-bromomethylacridine based on micellar phase-transfer catalysis
by F. A. L. van der Horst and M. H. Post (Utrecht, The Netherlands) and J. J. M. Holthuis and U. A. Th. Brinkman (Amsterdam, The Netherlands) (Received June 26th, 1989) . . . 443

On-line combination of dialysis and column-switching liquid chromatography as a fully automated sample preparation technique for biological samples. Determination of nitrofuran residues in edible products
by M. M. L. Aerts and W. M. J. Beek (Wageningen, The Netherlands) and U. A. Th. Brinkman (Amsterdam, The Netherlands) (Received October 13th, 1989) 453

Separation of ionic drug substances by supercritical fluid chromatography
by W. Steuer, J. Bauman and F. Erni (Basle, Switzerland) (Received September 25th, 1989) 469

Intrinsic fluorescence studies of the kinetic mechanism of unfolding of α-lactalbumin on weakly hydrophobic chromatographic surfaces
by P. Oroszlan, R. Blanco, X.-M. Lu, D. Yarmusch and B. L. Karger (Boston, MA, U.S.A.) (Received November 1st, 1989) 481

Microenvironmental contributions to the chromatographic behavior of subtilisin in hydrophobic-interaction and reversed-phase chromatography
by R. M. Chicz and F. E. Regnier (West Lafayette, IN, U.S.A.) (Received August 29th, 1989) 503

Reversed-phase high-performance liquid chromatography of proteins and polypeptides on polystyrene-coated silica supports
by V. A. Davankov and A. A. Kurganov (Moscow, U.S.S.R.) and K. K. Unger (Mainz, F.R.G.) (Received October 13th, 1989) 519

Differential interaction of peptides and protein surface structures with free metal ions and surface-immobilized metal ions
by T. W. Hutchens and T.-T. Yip (Houston, TX, U.S.A.) (Received October 5th, 1989) . 531

High-performance liquid chromatography of proteins on deformed non-porous agarose beads. Fast boronate affinity chromatography of haemoglobin at neutral pH
by S. Hjertén and J.-P. Li (Uppsala, Sweden) (Received October 26th, 1989) 543

Recent developments in the chromatographic analysis of carbohydrates (Review)
by S. C. Churms (Cape Town, South Africa) (Received October 11th, 1989) 555

Carbohydrates as a tool for oriented immobilization of antigens and antibodies
by J.Turková, L. Petkov, J. Sjadok, J. Káš and M. J. Beneš (Prague, Czechoslovakia) (Received September 5th, 1989) 585

Application of high-performance hydrophobic-interaction chromatography to the characterization of recombinant DNA-derived human growth hormone
by S.-L. Wu and W. S. Hancock (South San Francisco, CA, U.S.A.) and B. Pavlu and P. Gellerfors (Stockholm, Sweden) (Received October 16th, 1989) 595

Liquid column chromatography of oxo compounds as their 2,4-dinitrophenylhydrazones with special reference to glucose–glycine Maillard reaction products
by I. Misik, R. Struzinsky, K. Macek and Z. Deyl (Prague, Czechoslovakia) (Received November 6th, 1989) 607

Hippuric acid and 3-carboxy-4-methyl-5-propyl-2-furanpropionic acid in serum and urine. Analytical approaches and clinical relevance in kidney diseases
by H. M. Liebich, J. I. Bubeck, A. Pickert, G. Wahl and A. Scheiter (Tübingen, F.R.G.) (Received August 24th, 1989) 615

Enantioselective determination of metoprolol in plasma by liquid chromatography on a silica-bonded α_1-acid glycoprotein column
by B.-A. Persson, K. Balmér and P.-O. Lagerström (Mölndal, Sweden) and G. Schill (Uppsala, Sweden) (Received May 26th, 1989) 629

High-performance liquid chromatographic determination of etretinate and all-*trans*- and 13-*cis*-acitretin in human plasma
by A. P. De Leenheer, W. E. Lambert, J. P. De Bersaques and A. H. Kint (Ghent, Belgium) (Received July 5th, 1989) 637

Evaluation of photometric high-performance liquid chromatographic data for the determination of benflurone metabolites in biological materials with and without concomitant use of a standard
by I. M. Hais, M. Nobilis and E. Kvasničková (Hradec Králové, Czechoslovakia) (Received October 16th, 1989) 643

Determination of trichloroethanol, the active metabolite of chloral hydrate, in plasma by liquid chromatography
by R. N. Gupta (Hamilton, Canada) (Received July 5th, 1989) 655

Identification power of thin-layer chromatographic colour reactions and integration of colour codes in a database for computerized identification in systematic toxicological analysis
by R. A. de Zeeuw, D. T. Witte and J.-P. Franke (Groningen, The Netherlands) (Received July 18th, 1989) 661

New option in capillary zone electrophoresis. Use of a transient ionic matrix (dynamic pulse)
by P. Boček, M. Deml and J. Pospíchal (Brno, Czechoslovakia) (Received May 31st, 1989) 673

Preparative purification of human monoclonal antibody isoforms in a multi-compartment electrolyser with immobiline membranes
by P. G. Righetti (Milan, Italy), E. Wenisch, A. Jungbauer and H. Katinger (Vienna, Austria) and M. Faupel (Basle, Switzerland) (Received June 22nd, 1989) 681

Formation of a cysteine–acrylamide adduct in isoelectric focusing gels
by M. Chiari, A. Manzocchi and P. G. Righetti (Milan, Italy) (Received June 14th, 1989) 697

Author Index 705